


Innovation Pathways to Decarbonize Kiln and Furnace Technology


Maxwell Pisciotta^a, Hélène Pilorgé^a, Justine Davids^b, Zachary Sasso^b, Shelvey Swett^b, and Jennifer Wilcox^a

^aChemical and Biomolecular Engineering Department, University of Pennsylvania, ^bChemical and Biomolecular Engineering Department, Worcester Polytechnic Institute

Research Motivation





Industrial scale furnaces and kilns are used to heat raw materials to produce lucrative materials such as cement, lime, glass, and iron and steel. Industrial processes such as these make up nearly 22% of all US emissions, of which, cement, lime, glass, and iron and steel are responsible for a third. The CO₂ from these processes stem from raw material chemical reactions as well as the fossil fuel used to provide heat. To ensure demand for these materials can be met while preventing additional emissions, strategic innovation pathways need to be considered.




Natural Gas

Co-Location Opportunities







Results

Innovation Pathway	Rotary	Hearth	BF/BOF	EAF
CCS	+	+	-	-
Biomass Fuel Switching	++		-	N/A
Hydrogen Fuel Switching				N/A
Fluidized Bed	+	N/A	N/A	N/A
Oxyfuel with CCS	•	+	+	N/A
Electric	+			++

- ++ Industrial Scale
- In Development
- + Small Scale Research
- **---** Little Exploration

N/A: Not applicable pathway

In addition to the pathways discussed here, other technologies are being investigated to also meet the heating demands of these processes. These include:

- Direct Separation Calcination at Calix
- Steel production through electrolysis at Boston Metal
- Calcination using fuel cell waste heat at Origen Power

Current Projects & Future Work

Current Projects and Implementations:

- Biomass used in US Cement plants today: wood chips, pecan shells, rice hulls, pistachio shells, and peanuts.
- Oxyfuel is at industrial-scale in most of the glass industry
- LefargeHolcim oxyfuel calciner at Retznel Plant (Austria)
- Heidelberg oxyfuel calciner at Colleferro Plant (Italy)
- Al Reyadah Steel plant (Abu Dhabi, UAE)
- Minerals Technologies fluidized bed lime kiln (USA)

Conclusions and Future Work

The more mature technology, such as oxyfuel in the glass industry and CCS, need to deployed soon to avoid increasing atmospheric CO₂ emissions. However, further research and testing of high-grade heating options are needed to validate the use of the emerging technologies in various industries. This research can be facilitated through strategic partnerships between the industrial sector and energy or feedstock producers. These partnerships can then lead to mutually beneficial co-location to further realization of CO₂-reducing pathways. Co-location will lead to a reduction in transportation costs for feedstocks and byproducts, including minerals, oxygen, and CO₂ for storage or utilization.

Resources

attps://ghgdata.epa.gov/ghgp/main.do 2] Barker DJ, Turner SA, Napier-Moore PA, Clark M, Davison JE. CO2 Capture in the Cement Industry. Energy Procedia. 2009;1(1):87–94 3] Levine E, Jamison K. Oxy-Fuel Firing for the Glass Industry: An Update on the Impact of This Successful Government-Industry Cooperative Effort. In 2001. p. 9. 1] Gimenez M. CCUS Projects at LafargeHolcim Focus on Oxycombustion [Internet]. ECRA/CEMCAP Workshop; 2015 Sep 16 [cited 2020 Oct 9]; Dusseldorf. Available from: https://www.sintef.no/globalassets/sintef-energi/cemcap/gimenez ccus projects-at-lafargeholcimco2capture.pdf/ [5] von Scheele J. Oxyfuel Combustion in the Steel Industry: Energy Efficiency and Decrease of CO2 Emissions. In: Palm J, editor. Energy Efficiency [Internet]. Sciyo; 2010 [cited 2020] Oct 14]. Available from: http://www.intechopen.com/books/energy-efficiency/oxyfuel-combustion-in-the-steel-industry-energy-efficiency-and-decrease-of-co2-emissions [6] U.S. Geological Survey (USGS), 2003. Active mines and minerals plants in the US [WWW Document] (accessed 10.12.20) 9] U.S. Department of Agriculture (USDA), 2019. CropScape – Cropland Data Layer. National Agricultural Statistics Service. https://massgeodata.gmu.edu/CropScape 1] https://www.messergroup.com/documents/20182/1119059/Company+presentation+2019/7d4a19d3-d762-4cdd-Bdb0-19740aaa9d6c?version=1.18 [2] USGS, 2013. National Assessment of Geologic Carbon Dioxide Storage Resources, https://pubs.usgs.gov/ds/774/ Hartmann, J., Moosdorf, N., 2012. The new global lithological map database GLiM: A representation of rock properties at the Earth surface. Geochem. Geophys. Geosystems 13. 4] Whittaker, J.M., Afonso, J.C., Masterton, S., Muller, R.D., Wessel, P., Williams, S. E., Seton, M., 2015. Long-term interaction between mid-ocean ridges and mantle plumes. Nat. Geosci. 8, 479-483. https://doi.org/10.1038/ngeo2437 [7] Johansson, L., Zahirovic, S., Muller, R. D., 2018. The interplay between the eruption and weathering of large igneous provinces and the deep-time carbon cycle. Geophys. Res. Lett. 45, [6] Coleman R. G., Irwin W. P. (1974) Ophiolites and ancient continental margins. In: Burk, C. A., Drake C.L. (eds) The Geology of Continental Margins. Springer, Berlin, Heidelberg [7] Kirchofer, A., Brandt, A., Krevor, S., Prigiobbe, V., Becker, A. and Wilcox, J., 2013. Assessing the potential for mineral carbonation with industrial alkalinity sources in the US.

19] World Resources Institute (WRI), Global Power Plant Database, 2019, https://datasets.wri.org/dataset/globalpowerplantdatabas 20] Hart Energy Publishing, Carbon dioxide (CO₂) pipelines in the United States, 2014, http://purl.stanford.edu/yf510bb3744